



Corporate Presentation

AUGUST 2022

Forward Looking Statements

This presentation (together with any other statements or information that we may make in connection herewith) contains forward-looking statements within the meaning of the Private Securities Litigation Reform Act of 1995 with respect to Kiniksa Pharmaceuticals, Ltd. (and its consolidated subsidiaries, collectively, unless context otherwise requires, “Kiniksa,” “we,” “us” or “our”). In some cases, you can identify forward looking statements by terms such as “may,” “will,” “should,” “expect,” “plan,” “anticipate,” “could,” “intend,” “goal,” “design,” “target,” “project,” “contemplate,” “believe,” “estimate,” “predict,” “potential,” “strategy,” or “continue” or the negative of these terms or other similar expressions, although not all forward-looking statements contain these identifying words. All statements contained in this presentation that do not relate to matters of historical fact should be considered forward-looking statements, including without limitation, statements regarding our strategy; potential value drivers; potential indications; potential market opportunities and competitive position; ongoing, planned and potential clinical trials and other studies; timing and potential impact of clinical data; regulatory and other submissions, applications and approvals; commercial strategy and commercial activities; expected run rate for our cash, cash equivalents and short-term investments; expected funding of our operating plan; financial guidance; third-party collaborations and licensing; and capital allocation.

These statements involve known and unknown risks, uncertainties, and other important factors that may cause our actual results, performance or achievements to be materially different from those expressed or implied by the forward-looking statements, including, without limitation, potential delays or difficulties with our clinical trials; potential inability to demonstrate safety or efficacy or otherwise producing negative, inconclusive or uncompetitive results; potential for changes in final data from preliminary or interim data; potential inability to replicate in later clinical trials positive results from earlier trials and studies; our reliance on third parties for manufacturing and conducting clinical trials, research and other studies; our ability to realize value from our licensing and collaboration arrangements; our ability to source sufficient drug product, as needed, to meet our clinical and commercial requirements; our inability to demonstrate safety and efficacy to the satisfaction of applicable regulatory authorities; potential for applicable regulatory authorities to not accept our filings or to delay or deny approval of, or emergency use authorization for, any of our product candidates or to require additional data or trials to support any such approval or authorization; delays, difficulty or inability to successfully execute on our commercial strategy for ARCALYST; potential changes in our strategy, clinical trial priority, operating plan and funding requirements; raw materials, important ancillary product and drug substance and/or drug product shortages; substantial new or existing competition; potential impact of the COVID-19 pandemic or any subsequent pandemic, and measures taken in response to such pandemics, on our business and operations as well as the business and operations of our manufacturers, CROs upon whom we rely to conduct our clinical trials, and other third parties with whom we conduct business or otherwise engage, including the FDA and other regulatory authorities; risks related to the ongoing war in Ukraine; and our ability to attract and retain qualified personnel.


These and the important factors discussed in our filings with the U.S. Securities and Exchange Commission, including under the caption “Risk Factors” contained therein could cause actual results to differ materially from those indicated by the forward-looking statements made in this presentation. These forward-looking statements reflect various assumptions of Kiniksa's management that may or may not prove to be correct. No forward-looking statement is a guarantee of future results, performance, or achievements, and one should avoid placing undue reliance on such statements. Except as otherwise indicated, this presentation speaks as of the date of this presentation. We undertake no obligation to update any forward-looking statements, whether as a result of new information, future events or otherwise, except as required by law.

This presentation also contains estimates, projections, and/or other information regarding our industry, our business and the markets for certain of our product candidates, including data regarding the estimated size of those markets, and the incidence and prevalence of certain medical conditions. Unless otherwise expressly stated, we obtained this industry, business, market and other data from reports, research surveys, clinical trials, studies and similar data prepared by market research firms and other third parties, from industry, medical and general publications, and from government data and similar sources. Information that is based on estimates, forecasts, projections, market research, or similar methodologies is inherently subject to uncertainties and actual events or circumstances may differ materially from events and circumstances reflected in this information.

ARCALYST is a registered trademark of Regeneron Pharmaceuticals, Inc. All other trademarks are the property of their respective owners.



Portfolio of Immune-Modulating Assets



LICENSE AGREEMENT

Vixarelimab
OSMRβ

Roche and Genentech

GLOBAL RIGHTS FOR ALL INDICATIONS³



1) Approved in the U.S; 2) The FDA granted Breakthrough Therapy designation to ARCALYST for recurrent pericarditis in 2019; the FDA granted Orphan Drug exclusivity to ARCALYST in March 2021 for the treatment of recurrent pericarditis and reduction in risk of recurrence in adults and pediatric patients 12 years and older. The European Commission granted Orphan Drug designation to ARCALYST for the treatment of idiopathic pericarditis in 2020; 3) In August 2022, Kiniksa entered into an agreement to grant Genentech and Roche exclusive global rights to develop and commercialize vixarelimab, subject to certain closing conditions; 4) Phase 2 clinical trials of mavrilimumab in rheumatoid arthritis and giant cell arteritis achieved their primary and secondary endpoints with statistical significance; 5) Kiniksa granted Huadong Medicine exclusive rights in the Asia Pacific Region, excluding Japan; IL-1α = interleukin-1α ; IL-1β = interleukin-1β; GM-CSFRα = granulocyte macrophage colony stimulating factor receptor alpha; OSMRβ = oncostatin M receptor beta; MENA = Middle East and North Africa

Strategic Collaboration with Huadong Medicine for Asia Pacific Region

- Collaboration to develop and commercialize ARCALYST and mavrilimumab in the Asia Pacific Region (excluding Japan).
- Kiniksa received \$22 million upfront and is eligible to receive up to approximately \$640 million in specified development, regulatory and sales-based milestones.
 - The upfront payment included \$12 million for the territory license of ARCALYST and \$10 million for the territory license of mavrilimumab.
- Kiniksa could also receive tiered royalties ranging from the low-teens to the low-twenties on annual net sales.
- Kiniksa retains all existing development and commercialization rights for both assets outside of the Asia Pacific Region.

Accelerating our ability to bring multiple therapeutics to patients suffering from severe autoimmune and inflammatory diseases across the Asia Pacific Region

Collaboration provides non-dilutive capital and additional resources to help accelerate our development and commercialization efforts



Global License Agreement with Genentech for Vixarelimab

- Kiniksa will receive \$100 million in upfront and near-term payments, and is eligible to receive up to approximately \$600 million in certain clinical, regulatory, and sales-based milestones, before fulfilling upstream financial obligations.
- Kiniksa is also eligible to receive royalties on annual net sales. Genentech will obtain global rights for the development and commercialization of vixarelimab.
- Kiniksa has completed screening patients for the Phase 2b clinical trial of vixarelimab in prurigo nodularis. The company plans to complete the trial but will no longer disclose data in the second half of 2022.

\$100 million in non-dilutive proceeds from the transaction to advance synergistic cardiovascular opportunities



ARCALYST[®]

Arcalyst[®]
(rilonacept) For Injection

IL-1 α AND IL-1 β CYTOKINE TRAP

DISEASE AREA: Recurrent pericarditis¹; painful and debilitating auto-inflammatory cardiovascular disease

COMPETITION²: First and only FDA-approved therapy for recurrent pericarditis

REGULATORY: U.S. Orphan Drug exclusivity for treatment of and reduction in risk of recurrence of recurrent pericarditis; European Commission Orphan Drug designation in idiopathic pericarditis

STATUS: FDA-Approved

ECONOMICS: 50/50 profit split on profit and third-party proceeds

RIGHTS: Kiniksa has worldwide rights³ (excluding MENA) for all indications outside those in oncology and local administration to the eye or ear



1) ARCALYST is also approved and marketed for Cryopyrin-Associated Periodic Syndromes (CAPS) and maintenance of remission of Deficiency of Interleukin-1 Receptor Antagonist (DIRA) in the United States; 2) Drugs@FDA: ARCALYST Prescribing Information, Ilaris Prescribing Information, Kineret Prescribing Information; Kaiser et al. Rheumatol Int (2012) 32:295–299; Theodoropoulou et al. Pediatric Rheumatology 2015, 13(Suppl 1):P155; Fleischmann et al, 2017 ACR/ARHP Abstract 1196; Kosloski et al, J of Clin Pharm 2016, 56 (12) 1582-1590; Cohen et al. Arthritis Research & Therapy 2011, 13:R125; Cardiel et al. Arthritis Research & Therapy 2010, 12:R192; Hong et al. Lancet Oncol 2014, 15: 656-666; 3) Kiniksa granted Huadong Medicine exclusive rights in the Asia Pacific Region, excluding Japan; IL-1 α = interleukin-1 α ; IL-1 β = interleukin-1 β ; MENA = Middle East North Africa

ARCALYST Label

ARCALYST is a patient-administered once-weekly subcutaneous therapy

ADULTS (18 years and older)	ADOLESCENTS (12 to 17 years)
Loading dose: 320 mg delivered as two 160 mg (2 mL) injections	Loading dose: 4.4 mg/kg delivered up to a maximum of 320 mg, delivered as 1 or 2 injections (not to exceed 2 mL/injection)
Weekly maintenance dose: 160 mg delivered once weekly as a 2 mL injection	Weekly maintenance dose: 2.2 mg/kg delivered up to a maximum of 160 mg (2 mL) injection, once weekly


The first injection of ARCALYST should be performed under the supervision of a healthcare professional.



ARCALYST is supplied in sterile, single-use, 20-mL glass vials

- Each vial contains 220 mg ARCALYST, a sterile, white to off-white lyophilized powder
- Reconstitution with 2.3 mL of preservative-free Sterile Water for Injection is required prior to subcutaneous administration of the drug
- The reconstituted ARCALYST is a viscous, clear, colorless to pale yellow, free from particulates, 80-mg/mL preservative-free solution

Pericarditis Epidemiology




All figures annual period prevalence


Approximately 14,000 recurrent pericarditis patients in the U.S. suffer from persistent underlying disease, with multiple recurrences and inadequate response to conventional therapy¹

- **~ 160,000:** Epidemiological analysis using large national surveillance databases to calculate the pooled annualized prevalence of pericarditis (**Basis for Orphan Drug Designation approval**)²
- **~40,000:** Up to 30% experience at least one recurrence; some recur over multiple years^{6,7}
- **~14,000:** Nearly 50% annual turnover with ~7,000 patients coming into the pool each year⁸

Early Treated Patients Are Closely Associated to the Launch Target Population, While Prescribers Can Utilize ARCALYST Earlier in the Disease



ARCALYST PATIENTS BY PRIOR PRODUCT¹



ARCALYST PATIENTS BY FLARE STATUS @ INITIATION¹



Klein A, Cremer P, Kontzias A, Furqan M, Tubman R, Roy M, Magestro M. Annals of Epidemiology. 2019;36:71; 2) Lin D, Majeski C, DerSarkissian M, Magestro M, Cavanaugh C, Laliberte F, Lejune D, Mahendran M, Duh M, Klein A, Cremer P, Kontzias A, Furqan M, Tubman R, Roy M, Mage. (Nov, 2019). Real-World Clinical Characteristics and Recurrence Burden of Patients Diagnosed with Recurrent Pericarditis in the United States. Poster session presented at the American Heart Association, Philadelphia, PA.; 3) ClearView Forecasting Analysis 2019 Q1

Source: 1) Kiniksa Pharmaceuticals data on file 2021. 2) Other late line agents include anakinra, azathioprine, methotrexate

Collaborative Field Force to Drive Awareness, Overcome Access Barriers and Help Ensure Positive Patient and Physician Experience



SALES

CLINICAL SALES SPECIALISTS

- **Focus:** ~2500 HCPs across ~800 accounts
- **Responsibility:** Physician accounts, disease education, ARCALYST promotion, account and territory plans, speaker program planning

PAYER

STRATEGIC ACCOUNTS

- **Focus:** ~350 payers and 5 Specialty Pharmacies
- **Responsibility:** Payer/specialty pharmacy relationship, strategic account planning, support sales team

MEDICAL

MEDICAL SCIENCE LIAISONS

- **Focus:** Subject Matter Experts and HCPs
- **Responsibility:** Disease awareness, data dissemination, advocacy development, account and payer support, speaker management

PATIENT ACCESS

KINIKSA ONECONNECT™ PROGRAM

- **Focus:** Patients and caregivers, HCPs seeking reimbursement support for their patients
- **Responsibility:** Optimize patient and customer experience with ARCALYST and Kiniksa, provide seamless initiation, reimbursement, and adherence support



HCP = health care provider

Pricing, Access and Distribution Considerations

Pricing

- ARCALYST list price of \$20,700 per month
Based on first and only FDA-approved therapy for recurrent pericarditis, in-line with specialty biologics with Breakthrough Therapy and Orphan Drug designation
- Helping to ensure **patient affordability** and access to treatment is one of our core principles and to this end, we offer a suite of programs to support affordability to eligible patients who are prescribed ARCALYST

Access

- Kiniksa's goal is to enable rapid and broad access to ARCALYST for patients with Recurrent Pericarditis, CAPS, and DIRA
- Payer mix for ARCALYST is largely **commercial (~70%), Medicare (~20%), Medicaid (~10%)**
- Early payer engagement has increased awareness of recurrent pericarditis and the differentiated value of ARCALYST
- The **Kiniksa OneConnect™** program is a personalized treatment support program for patients prescribed ARCALYST

Distribution

- ARCALYST is distributed **through a closed network of designated specialty pharmacies and the Veterans Affairs**
- The distribution network for ARCALYST was developed to provide a high and consistent level of patient support with broad access. Network pharmacies provide customized services to support patients

Continued ARCALYST Growth in Q2 2022; On-track for Annual Guidance of \$115-130M Net Revenue

Net Revenue

- ARCALYST net revenue for Q2 2022 was \$27.0 million
- Represents +22% net revenue growth versus Q1 2022
- Represents +250% net revenue growth versus Q2 2021

*\$27.0 million in net revenue for the **second quarter of 2022***

*\$49.2 million net revenue for the **first half of 2022***

Revenue Drivers

- Strong recurrent pericarditis demand was the primary growth lever, driven by new patient initiations
- CAPS and DIRA patient demand remained stable and broadly consistent with the previous quarters
- Growth rate represents continued uptake and adoption of ARCALYST from prescribers, payers and patients in this previously unmet and debilitating autoinflammatory cardiovascular disease

*\$80.0 million net revenue for the most recent **12-months***

2022 Guidance of \$115-130 million net revenue



Drivers of Continued ARCALYST Growth in Recurrent Pericarditis

Physician Growth

- Continued growth in prescriber base with more than 550 HCPs having prescribed ARCALYST since launch
- Growing repeat prescriber base with 20% having prescribed ARCALYST for 2 or more patients

Payer Access

- Payers continue to recognize value of the first and only approved drug for recurrent pericarditis, with greater than 90% approval rate of completed cases


Duration of Therapy

- ARCALYST use in recurrent pericarditis to date suggest an average continuous treatment durations of approximately 12 months
 - Majority of prescribers writing for 12 months
 - Majority of payers approved for 12 months prior to re-authorization
 - Adherence and compliance continues to be strong
 - Approximately ~45% of Q2 2021 initiations remain on continuous therapy at end of Q2 2022

Patient Satisfaction¹

- Patients report high satisfaction when treated with ARCALYST, especially when compared to other conventional therapeutic options

Total and Repeat Prescriber Growth per Quarter



■ Prescribers with ≥2 Recurrent Pericarditis Enrollments
■ Prescribers with 1 Recurrent Pericarditis Enrollment



1: Among Patients with prior experience on each therapy for recurrent pericarditis. Data on file.

HCPs: Healthcare Professionals

Field Force Expansion to Accelerate Penetration of US Addressable Market


Field Force Effectiveness

- Physicians who meet with our field team have a significantly higher awareness of recurrent pericarditis and ARCALYST
- Additionally, the intention to prescribe ARCALYST also increases significantly


Opportunity in our Target Population

- Targeting the annual target pool of 14K patients with 2 or more recurrences, acknowledging ARCALYST’s broad label could grow the pool in appropriate patients on the first recurrence
- With only a few specialist pericardial disease centers, patients are widely dissipated across the US and often present to different cardiologists even in the same institution compared to their prior flare(s)
- Frequent call activity is a major driver of prescriber understanding of disease burden and ARCALYST adoption
- Growing confidence in our launch trajectory and a profitable collaboration allows us to expand our field team to enable us to reach patients quicker

EXPECTATION OF ARCALYST PRESCRIBING OVER NEXT 6 MONTHS¹



14,000 TARGET PATIENTS WITH UPSIDE IN FIRST RECURRENCE POOL



All figures annual period prevalence



1: Among Cardiologists. Data on file.

Field Evolution to Create Greater Reach and Frequency with Top Tier Doctors as well as Reach to a Broader Set of Physicians

Field Launch Strategy

LEAN TEAM WITH FOCUSED &
TARGETED EXECUTION

~30 Specialty Cardiology Reps

Initial launch focus on
top tier accounts:

~3,300 individual prescribers

~45% of RP patients nationally



Following adoption, moving into next deciles

Strategy Evolution

EXPANDED TEAM CREATING
GREATER REACH AND FREQUENCY

~50 Specialty Cardiology Reps

Increased focus within top tier accounts as
well as expanded reach at mid tier
prescribers, reaching:

~6,000 top and mid tier prescribers

~70% of RP patients nationally

Plus, supplemented by a small Inside Sales
Team as a cost-efficient approach to
reaching lower decile physicians



Evolved field team will be in place in Q4 2022

Expanded Live Education and Sales Engagements Anticipated to Drive Further Adoption

Increasing levels of prescriber awareness with more significant opportunities to further broaden knowledge on horizon

- Return of live national and regional conferences after pandemic move to virtual
- Significant attendee engagement at American College of Cardiology (ACC), held in April '22, though booth traffic and product theatre
- American College of Rheumatology (ACR) and American Heart Association (AHA) coming in November 2022

Sales interactions are having a major impact with target base

- Prescribers with recent field sales representative interactions are more knowledgeable and subsequently indicate higher expectation of increasing ARCALYST use

Most new ARCALYST patients continue to come from existing field targets

- Additional prescribing also coming from the long tail of prescribers treating 1-2 recurrent pericarditis patients per year

ACC HIGHLIGHTS




ARCALYST BOOTH
>400 individual details with
Cardiologists



ARCALYST PRODUCT THEATRE
>120 prescribers attended our
branded symposium

EXPECTATION OF ARCALYST PRESCRIBING OVER NEXT 6 MONTHS¹



■ Unaware of Arcalyst ■ Decrease ■ Stay About the Same ■ Increase




1: Among Cardiologists. Data on file.

Aligned with our Data and Promotional Campaign, we are Seeing Momentum to Move ARCALYST Ahead of, or as an Alternative to, Corticosteroids

RECURRENT PERICARDITIS THOUGHT LEADER POSITIONING OF ARCALYST IN TREATMENT ALGORITHM¹

Acute Pericarditis	Index Episode	• NSAIDs (week(s)) • Colchicine (3 months)
Recurrent Pericarditis	First Recurrence	• NSAIDs (weeks to months) • Colchicine (≥6 months)
	Second (or more) Recurrence	• ARCALYST (rilonacept), including ahead of corticosteroids and other drugs if intolerant
Additionally	• ARCALYST can be used in colchicine resistant and/or corticosteroid dependent patients, including while tapering down steroids and other agents	

ARCALYST PATIENTS BY PRIOR PRODUCT²



MINIMAL STEP EDITS FOR ARCALYST¹




% of covered lives



1) Wang and Klein. Current Cardiology Reports (2022) 24:23-30. <https://doi.org/10.1007/s11886-021-01621-0>; 2) Kiniksa Pharmaceuticals data on file 2022. 3) Other late line agents include anakinra, azathioprine, methotrexate

Summary of ARCALYST Profit Share Arrangement with Regeneron¹



- Kiniksa is responsible for sales and distribution of ARCALYST in all approved indications in the United States.
- Kiniksa's license to ARCALYST includes worldwide rights, excluding the Middle East and North Africa, for all applications other than those in oncology and local administration to the eye or ear.
- Kiniksa covers 100% of development expenses related to approval of additional indications.
- We evenly split profits on sales with Regeneron



1) Subject to description contained in definitive agreement; 2) Global net sales for CAPS, DIRA and recurrent pericarditis recognized as revenue on Kiniksa's income statement; 3) Including cost of product purchased from Regeneron as well as relevant Kiniksa overhead; CAPS = Cryopyrin-Associated Periodic Syndromes; DIRA = Deficiency of the Interleukin-1 Receptor Antagonist

KPL-404

MONOCLONAL ANTIBODY INHIBITOR INTERACTION BETWEEN CD40 AND CD154

DISEASE AREA: Rheumatoid Arthritis; a chronic inflammatory disorder affecting many joints; External proof-of-concept previously established in broad range of autoimmune diseases: Sjogren's disease, systemic lupus, solid organ transplant and Graves' disease¹

SCIENTIFIC RATIONALE^{2,3}: Attractive target for blocking T-cell dependent, B-cell-mediated autoimmunity

STATUS: Phase 1 single-ascending-dose study in healthy volunteers completed and supports further development in patients with optionality for testing SC and/or IV dosing

ECONOMICS: Negligible clinical and regulatory milestones and royalty on annual net sales


RIGHTS: Worldwide



1) Poster presentation at the Keystone Symposia: Antibodies as Drugs: New Horizons in the Therapeutic Use of Engineered Antibodies: KPL-404, a CD40 antagonist, blocked antigen-specific antibody responses in an in vivo NHP model and demonstrated strong PK/PD correlation; 2) Elgueta, et al. Immunol Rev 2009, 229 (1), 152-172; 3) Peters, et al. Semin Immunol 2009, 21 (5) 293-300; RO = receptor occupancy; TDAR = T-cell Dependent Antibody Response


CD40/CD154 is an Essential Immune Pathway for T-Cell Priming and T-Cell Dependent B-Cell Responses

- CD40 is expressed on the surface of dendritic cells, B-cells, antigen-presenting cells and non-immune cell types
- Its ligand, CD40L (CD154), is expressed by activated T-cells, platelets, and other cell types



- CD40 ligation on DCs induces cell maturation by promoting antigen presentation and enhancing their costimulatory activity
- Mature DCs stimulate activated T-cells to increase IL-2 production that facilitates T-helper cells (Th) and cytolytic T-Lymphocyte (CTL) expansion
- CD40-stimulated DCs also secrete cytokines favoring Th1 cell differentiation and promoting Th cell migration to sites of inflammation
- CD40 ligation also provides a pro-inflammatory signal within the mononuclear phagocyte system


- Humoral immunity is dependent on a thriving B cell population and activation by Th cells; blockade of CD40/CD40L interaction has been shown to completely ablate primary and secondary TDAR response



- CD40 engagement triggers B-cell intercellular adhesion, sustained proliferation, expansion, differentiation, and antibody isotype switching leading to affinity maturation, which is essential for generation of memory B cells and long-lived plasma cells

- B-cells require contact-dependent stimulus from T cells through CD40/CD40L interaction independent of cytokines to trigger growth and differentiation

KPL-404 Single-Ascending-Dose Phase 1 Study



Notes: Unless otherwise noted dose groups included 6 active/2 placebo subjects; *1° KLH challenge for all SAD dose groups except 0.03 and 0.3 mg/kg, 2° KLH re-challenge only in 1, 3, and 10 mg/kg IV; ** Cohort included 2 active and 2 placebo subjects

Final Data from KPL-404 Single-Ascending-Dose Phase 1 Study

The randomized, double-blind, placebo-controlled first-in-human (FIH) study is designed to investigate the safety, tolerability, PK and PD properties of single-ascending intravenous (IV) and subcutaneous (SC) doses of KPL-404 in healthy subjects.

- 2 single-ascending-dose arms (SAD):
 - Single-dose KPL-404 0.03 mg/kg, 0.3 mg/kg, 1 mg/kg, 3 mg/kg or 10 mg/kg IV and
 - Single-dose KPL-404 1 mg/kg or 5 mg/kg SC

Primary Endpoint: Safety and tolerability of single ascending intravenous (IV) and subcutaneous (SC) doses of KPL-404 in healthy subjects.

- KLH challenge in 1 mg/kg, 3 mg/kg, and 10 mg/kg IV and 1 mg/kg and 5 mg/kg SC cohort

Secondary Endpoints: Pharmacokinetics and anti-drug antibody response following single IV and SC doses of KPL-404 in healthy subjects, serum anti- keyhole limpet hemocyanin (KLH) IgG levels

Exploratory Endpoint: Receptor occupancy of KPL-404 on CD40 in healthy subjects

Preliminary Data:


- All dose escalations occurred as per protocol with no dose limiting safety findings. All 6 subjects dosed with KPL-404 3 mg/kg IV showed full receptor occupancy through Day 29, which corresponded with complete suppression of the T-cell Dependent Antibody Response (TDAR) to KLH through Day 29. Consistent dose relatedness was shown in the lower dose level cohorts, including 0.03 mg/kg, 0.3 mg/kg, 1 mg/kg IV and 1 mg/kg SC. Data collection for the higher dose level cohorts, 10 mg/kg IV and 5 mg/kg SC, is ongoing.
- The data to-date support subsequent study in patients, including potential IV or SC monthly administration.

Final Data:

- KPL-404 showed dose-dependent increases in concentration across cohorts. All dose escalations occurred as per protocol with no dose-limiting safety findings.
- KPL-404 was well-tolerated, and there were no serious adverse events.
- Subjects dosed with KPL-404 10 mg/kg IV showed full RO through at least Day 71 and complete suppression of TDAR after KLH challenge and re-challenge through at least Day 57.
- Subjects dosed with KPL-404 5 mg/kg SC showed full RO through Day 43 and suppression of TDAR after KLH challenge through at least Day 29. These data confirm and extend previously-reported 3 mg/kg IV cohort data, in which RO and suppression of TDAR after KLH challenge were demonstrated through Day 29.
- The 3 mg/kg IV dose level had previously demonstrated complete suppression of memory TDAR response to a re-challenge on Day 29.
- Anti-drug antibodies to KPL-404 were suppressed for at least 57 days at 10 mg/kg IV; the suppression of antibody responses to the drug itself is an independent indicator of target engagement and pharmacodynamic effect


Final Data from KPL-404 Single-Ascending-Dose Phase 1 Study

Pharmacokinetic profiles for KPL-404



Final Data from KPL-404 Single-Ascending-Dose Phase 1 Study

T-Cell Dependent Antibody Response (TDAR) for KLH antigen challenge



KLH = keyhole limpet hemocyanin

KPL-404 Phase 2 Trial in Rheumatoid Arthritis Ongoing

12-week study designed to provide PK data and early signal of efficacy with chronic administration, and optionality to evaluate KPL-404 across a range of other autoimmune diseases


Cohorts 1 through 3 will each sequentially randomize eight patients
Cohort 4 will randomize up to 60 patients

PATIENT POPULATION :

- Active RA who have an inadequate response to or are intolerant to a Janus kinase inhibitor (JAKi) or at least one biologic disease-modifying anti-rheumatic drug (bDMARD). Subjects who have failed both bDMARD and JAKi are excluded from the study.

DISEASE CRITERIA:

- Six or more swollen joints and ≥ 6 tender joints at screening and baseline line visits; levels of high sensitivity C-reactive protein ≥ 5 mg/L; seropositivity for serum RF and/or ACPA at screening.



COHORTS 1-3

- Each cohort will sequentially randomize 8 patients
- Primary Endpoints:
 - Incidence of treatment-emergent adverse events (TEAEs)
 - Pharmacokinetics (C_{max} , $AUC_{(0-t)}$)
- Secondary Endpoint:
 - Change from baseline in DAS28-CRP at Week 12

COHORT 4

- Cohort 4 will randomize up to 60 patients
- Primary Endpoint:
 - Change from baseline in DAS28-CRP at Week 12
- Secondary Endpoints :
 - Incidence of treatment-emergent adverse events (TEAEs)
 - Pharmacokinetics (C_{max} , $AUC_{(0-t)}$)

Objectives: Evaluate safety, efficacy, and PD compared with placebo across the estimated therapeutic range and to characterize PK across varying dose levels of KPL-404.


The protocol allows for additional cohorts to be initiated at a dose & regimen to be determined.




SC = subcutaneous; q2wk = every other week; AUC = Area Under the Curve; RF = Rheumatoid Factor; ACPA = anti-citrullinated protein antibodies, PD = Pharmacodynamics; PK = Pharmacokinetics; R = Randomization; SRC = Safety Review Committee

Potential for Evaluation of KPL-404 in a Broad Range of Autoimmune Diseases

Indications with Published Data



Indications with Pending Data & Trials Ongoing



*Indications evaluated with subcutaneous administration
1) With the CD40 mechanism

INDICATION SELECTION CRITERIA

- Robust Data or proof-of-concept supporting mechanism
- Differentiation vs. Competitors
- Commercial Attractiveness



Sources: 2019 numbers: <https://unos.org/data/transplant-trends/>; Hunter et al. Prevalence of rheumatoid arthritis in the United States adult population in healthcare claims databases, 2004-2014; Rheumatol Int. 2017 Sep;37(9):1551-1557; Overall Prevalence: Maciel et al, Arthritis Care Res (Hoboken) 2017; Qin et al, Ann Rheum Dis 2015; UpToDate; Baldini et al. Prevalence of Severe Extra-Glandular Manifestations in a Large Cohort of Patients with Primary Sjögren's Syndrome; 2012 ACR/ARHP Annual Meeting, ABSTRACT NUMBER: 2185; Wallin et al. The prevalence of MS in the United States A population-based estimate using health claims data, Neurology, March 5, 2019; Somers et al.; Prevalence of Systemic Lupus Erythematosus in the United States: Preliminary Estimates from a Meta-Analysis of the Centers for Disease Control and Prevention Lupus Registries; 2019 ACR/ARP Annual Meeting ABSTRACT NUMBER: 2886; Garg et al. JAMA Dermatol. 2017;153(8):760-764. doi:10.1001/jamadermatol.2017.0201 Sex- and Age-Adjusted Population Analysis of Prevalence Estimates for Hidradenitis Suppurativa in the United States; MayoClinic.org; Yale J Biol Med. 2013 Jun; 86(2): 255-260. N Engl J Med 2016;375:2570-81; <https://www.diabetesresearch.org/diabetes-statistics>; Nephcare.org; Kitiyakara C, Eggers P, Kopp JB. Twenty-one-year trend in ESRD due to focal segmental glomerulosclerosis in the United States. Am J Kidney Dis. 2004 Nov;44(5):815-25; Rachakonda et al. J Am Acad Dermatol. 2014 Mar;70(3):512-6. doi: 10.1016/j.jaad.2013.11.013. Epub 2014 Jan 2. Psoriasis prevalence among adults in the United States; Yeung et al. Psoriasis severity and the prevalence of major medical co-morbidities: a population-based study; JAMA Dermatol. 2013 Oct 1; 149(10): 1173-1179; Hoover et al. Kidney Int. 2016 Sep; 90(3): 487-492. Insights into the Epidemiology and Management of Lupus Nephritis from the U.S. Rheumatologist's Perspective.



Financials

Second Quarter 2022

Second Quarter 2022 Financial Results

Income Statement	Q2 2022	Q2 2021
Product Revenue	\$27.0M	\$7.7M
Collaboration Revenue	\$0.0M	\$0.0M
Total Revenue	\$27.0M	\$7.7M
Cost of Goods Sold	\$5.0M	\$2.5M
Collaboration Expenses	\$3.7M	\$0.0M
Research and Development Expenses	\$13.8M	\$23.9M
Selling, General and Administrative Expenses	\$23.8M	\$21.8M
Total Operating Expenses	\$46.3M	\$48.3M
Net Loss	(\$20.0M)	(\$41.6M)
Balance Sheet	June 30, 2022	December 31, 2021
Cash, Cash Equivalents and Short-term Investments	\$138.2M	\$182.2M

Cash reserves after the close of the vixarelimab global license agreement expected to fund operations into at least 2025

Second Quarter 2022 Collaboration Expense¹

ARCALYST Net Sales (RP + CAPS + DIRA)	\$27.0M	<i>Recognized as revenue on Kiniksa's income statement</i>
Cost of Goods Sold Related to Product Sales	(\$4.8M)	<i>Costs of product purchased as well as relevant overhead; amortization of ARCALYST commercial milestone excluded</i>
Commercial, Marketing, Regulatory and Other Expenses	(\$14.9M)	<i>100% of field force expense as well as commercial and marketing expenses subject to specified limits</i>
ARCALYST Operating Profit	\$7.4M	
Collaboration Expense	\$3.7M	<i>50% of ARCALYST operating profit booked as a separate line item within operating expenses</i>




1) Subject to the terms of the definitive agreements between Kiniksa and Regeneron; RP = Recurrent Pericarditis, CAPS = Cryopyrin-Associated Periodic Syndromes, DIRA = Deficiency of Interleukin-1 Receptor Agonist



Appendix

ARCALYST (rilonacept)


Role of IL-1 α and IL-1 β in the Autoinflammatory Cycle of Recurrent Pericarditis



Brucato A, et al. Int Emerg Med 2018 <https://doi.org/10.1007/s11739-018-1907-x>
Dinarello CA, et al. Nat Rev Drug Discov 2012;11:633-652

Addressable U.S. Opportunity of ARCALYST Estimated to be ~14K Patients

~7K new patients with multiple recurrences enter target pool annually



- ~7K new patients with repeat recurrences annually
- ~14K total patients with repeat recurrences annually at any point


Year	-4	-3	-2	-1	0
Incident case of acute pericarditis (1 st episode) ¹	117K	117K	117K	117K	117K
Incidence of initial RP patients (1 st recurrence) ²	26K	26K	26K	26K	26K
Ongoing recurrent from year-1 ³				7K	7K
Ongoing recurrent from year-2 ³			7K	3.5K	3.5K
Ongoing recurrent from year-3 ³		7K	3.5K	1.8K	1.8K
Ongoing recurrent from year-4 ³	7K	3.5K	1.8K	0.9K	0.9K
Ongoing recurrent from year-5 ³	3.5K	1.8K	0.9K	0.5K	0.5K
Ongoing recurrent from year-6 ³	1.8K	0.9K	0.5K	0.2K	0.2K
Ongoing recurrent from year-7 ³					0.1K

Addressable Opportunity in U.S.



1: Prevalence estimate from Imazio, et al. (2008); includes all etiologies (~80% idiopathic)
 2: Mid point of 15-30% of initial recurrence rate published in ESC Guidelines given higher colchicine use today
 3: Estimate for recurrence rate of subsequent recurrences from ESC Guidelines and Claims Analysis

Pivotal Phase 3 Trial of ARCALYST in Recurrent Pericarditis



Inclusion Criteria:

- All etiologies except infection and malignancy
- Present at screening with at least a third pericarditis episode, defined as at least 1 day with **NRS pain of ≥ 4** and **CRP value ≥ 1 mg/dL** within the 7-day period prior to first study drug administration
- Concomitant NSAIDs and/or colchicine and/or oral corticosteroid treatment in any combination

Primary Efficacy Endpoint :

- Time-to-first-adjudicated pericarditis-recurrence in the RW period

Major Secondary Efficacy Endpoints (16-weeks):

- Proportion of subjects who maintained Clinical Response
- Percentage of days with no or minimal pain
- Proportion of subjects with absent or minimal pericarditis symptoms


CEC Adjudication Criteria:

- Typical pericarditis pain (≥ 1 pain **NRS recording ≥ 4**) AND elevated **CRP (≥ 1.0 mg/dL)**, same day or ≤ 7 days
- Typical pericarditis pain (≥ 1 pain **NRS recording ≥ 4**) AND abnormal **CRP (>0.5 mg/dL)**, same day or ≤ 7 days AND ≥ 1 **supportive evidence** of pericarditis
- Typical pericarditis pain (BUT pain **NRS recording ≤ 4**) AND elevated **CRP (≥ 1.0 mg/dL)**, AND ≥ 1 **supportive evidence** of pericarditis

Baseline Demographics and Clinical Characteristics


Pivotal Phase 3 Riloncept Data

Baseline Demographics (n=86)




Total Number of Episodes Including Index and Qualifying Episodes	Run-in Period (n=86)
Mean	4.7


Prior Pericarditis History at Baseline (n=86)




SoC Received at Qualifying Episode (n=86)



Qualifying Episode CRP & NRS (n=86)




Pericarditis Manifestations at Qualifying Episode (n=86)



Subject Disposition

Pivotal Phase 3 Rilonacept Data

Median time to Monotherapy rilonacept = 7.9 weeks (7.0, 8.1)
 Median time to Pain Response² = 5.0 days (4.0, 6.0)
 Median time to CRP Normalization³ = 7.0 days (5.0, 8.0)
 Median time to Treatment Response⁴ = 5.0 days (4.0, 7.0)



1) Adverse Events n = 4 (4.7%); Protocol Deviation /Withdrawn Consent / Sponsor/Investigator Decision n = 3 (3.5%); CRP = C-reactive protein; NRS = Numerical Rating Scale 2; Pain response is NRS ≤ 2; 3) CRP normalization is CRP ≤ 0.5; 4) Treatment response is the combination of pain response and CRP normalization

ARCALYST Initiation Resulted in Rapid Resolution of Pericarditis Episodes

Pivotal Phase 3 RHAPSODY Data

Rapid and sustained reductions in both reported pain and inflammation as early as after the first dose of ARCALYST

Median time to pain response = 5.0 days; Median time to CRP normalization = 7.0 days

Secondary endpoints that were assessed during the run-in period

5 days

Time to treatment response
(median; 95% CI: 4, 7)*

97%

Treatment response* rate

7.9 weeks

Time to ARCALYST monotherapy
(median; 95% CI: 7, 8)



*Time to treatment response was defined as the time from the first dose to the first day when pericardial pain was NRS ≤ 2 and CRP ≤ 0.5 mg/dL (measured within 7 days before or after the pain response). During the 12-week run-in period, 77 of 79 patients demonstrated a treatment response.

Klein AL, Imazio M, Cremer P, et al. Phase 3 trial of interleukin-1 trap rilonacept in recurrent pericarditis. N Engl J Med. 2021;384(1):31-41.
ARCALYST (rilonacept) prescribing information 2021

ARCALYST Demonstrated a Steroid-Sparing Treatment Effect

Pivotal Phase 3 RHAPSODY Data

Patients treated with ARCALYST discontinued corticosteroids

In the run-in period of the Phase 3 trial RHAPSODY, patients receiving corticosteroids at baseline were transitioned to ARCALYST monotherapy in 7.9 weeks

Each patient treated with corticosteroids at baseline achieved clinical response with ARCALYST monotherapy

- 49% (27 of 86) of patients received corticosteroids at baseline
- None of the patients treated with corticosteroids at baseline and randomized to ARCALYST monotherapy experienced a recurrence while on therapy




96% Reduction in Risk of Pericarditis Recurrence

Pivotal Phase 3 RHAPSODY Data

ARCALYST reduced the risk of pericarditis recurrence

The primary efficacy endpoint was time to first adjudicated pericarditis recurrence in the randomized withdrawal period.



96%

reduction in the risk of recurrent pericarditis
(hazard ratio: 0.04; $p < 0.0001$)

The median time to recurrence on ARCALYST could not be estimated due to the low number of recurrences

- 2 of 30 of patients treated with ARCALYST had a recurrence
- The 2 pericarditis recurrences with ARCALYST occurred during temporary interruptions of 1 to 3 doses of ARCALYST

The median time to recurrence on placebo was 8.6 weeks (95% CI: 4.0, 11.7)

- 74% (23 of 31) of patients treated with placebo experienced a recurrence at the time that the event-driven portion of the trial was closed
- Consistent with the expected washout pharmacokinetics of once-weekly ARCALYST at steady state

92% of Trial Days of No/Minimal Pain

Pivotal Phase 3 RHAPSODY Data

Patients on ARCALYST had significantly more trial days with no/minimal pain vs placebo

Secondary efficacy endpoint was assessed during the randomized withdrawal period

92% of days

Patients reported no/minimal (NRS≤2) pericarditis pain

Compared with 40% of trial days in patients on placebo ($p<0.0001$) at the secondary endpoint assessed at Week 16 of the randomized withdrawal period.

At Week 16 of the randomized withdrawal period:

- A majority (81%) of patients maintained a clinical response measured at Week 16 of the randomized withdrawal period compared with 20% of patients on placebo ($p=0.0002$)



Most Common ARCALYST Adverse Reactions:

Injection-site reactions and upper respiratory tract infections

Adverse experiences in RHAPSODY

EVENT	RUN-IN PERIOD	RANDOMIZED-WITHDRAWAL PERIOD				TOTAL (N=86)
	Rilonacept (N=86)	Rilonacept, Including Bailout (N=30)	Placebo, Including Bailout (N=31) <i>number of patients with event (percent)</i>	Rilonacept, Before Bailout (N=30)	Placebo, Before Bailout (N=31)	
Any adverse event	69 (80)	24 (80)	22 (71)	24 (80)	13 (42)	74 (86)
Adverse events according to maximum severity [†]						
Mild	52 (60)	16 (53)	17 (55)	16 (53)	9 (29)	47 (55)
Moderate	15 (17)	8 (27)	5 (16)	8 (27)	4 (13)	25 (29)
Severe	2 (2)	0	0	0	0	2 (2)
Serious adverse event	1 (1)	1 (3)	3 (10)	1 (3)	1 (3)	5 (6)
Adverse event leading to death	0	0	0	0	0	0
Adverse event leading to dose interruption	0	1 (3)	0	1 (3)	0	1 (1)
Adverse event leading to discontinuation of rilonacept or placebo	4 (5)	0	0	0	0	4 (5)
Cancer [‡]	0	1 (3)	0	1 (3)	0	1 (1)
Injection-site reaction	28 (33)	6 (20)	2 (6)	5 (17)	0	29 (34)
Infection or infestation	14 (16)	12 (40)	7 (23)	12 (40)	3 (10)	29 (34)
Upper respiratory tract infection	12 (14)	7 (23)	2 (6)	7 (23)	0	19 (22)

*Patients with multiple events were counted once in each appropriate category

[†]Counted once, according to the maximum severity of the adverse event.

[‡]Cancer was an event of special interest.







Appendix KPL-404

KPL-404 Showed Encouraging Results in a Non-Human Primate Model of TDAR


Mean KPL-404 PK
(Intravenous Dosing)



Mean KPL-404 Receptor Occupancy (RO)



Mean KLH IgG



Showed linear pharmacokinetic profile with low variability between non-human primate subjects (n=7)

KPL-404 achieved 100% receptor occupancy for 2 weeks in all animals at 5mg/kg and 4 weeks in all animals at 10mg/kg

Complete suppression of primary T-cell dependent antigen response correlated with 100% receptor occupancy



Corporate Presentation

AUGUST 2022